


## Metolachlor-ESA

Investigating the transport and fate in soil and groundwater of a prominent metabolite of the corn herbicide s-metolachlor.



### Inspiration

The herbicide S-metolachlor has been increasingly applied to corn crops since 2005, when a ban was placed on the previously used herbicide, atrazine. Metolachlor-ESA, a major metabolite of s-metolachlor, has begun to appear in worrying concentrations of up to 1000 ng/l in tapped springs of the Luxembourg Sandstone aquifer. Because of the hydraulic inertia of the aquifer (mean residence times of ten years and longer) and its importance as a fresh water reservoir for Luxembourg, rapid action is necessary to determine whether contamination by metolachlor-ESA is due to the temporary activation of preferential flow paths after storm events and restricted to “reactive” springs, or is a warning sign indicating the long-lasting pollution of the majority of the springs draining the Luxembourg Sandstone resulting from the change in pesticide application practices.

### Innovation

The project will investigate the transport of metolachlor-ESA in the soil and in the aquifer of the Luxembourg Sandstone using a combination of field surveys, laboratory experiments and numerical simulations. Because soil and aquifer compartments are both major controls on the formation and transport of metolachlor-ESA, an integrated approach is necessary to predict future trends for different scenarios ranging from business-as-usual to an immediate ban of s-metholachlor.

Twelve springs covering a large range of metolachlor-ESA concentrations will be sampled weekly to characterize the pesticide dynamics and determine the fast-flow contribution to total discharge (both in terms of water and pesticide fluxes). Microcosm experiments will establish the transformation of metolachlor-ESA in soils under variable boundary conditions. Long-term soil leaching for different management scenarios will be simulated with the pesticide fate model PEARL calibrated using the data from the laboratory experiments and the spring monitoring. In a final step, the evolution of metolachlor-ESA concentrations in spring water will be predicted for each management scenario using a simple transit time model.

### Impact

The project will:

- quantify the interplay between degradation kinetics in soils and fast-flow dynamics
- identify Luxembourgish springs vulnerable to agricultural pollutants using metolachlor-ESA as a tracer

The dynamics of metolachlor-ESA in the springs will be unraveled using a combination of monitoring and modelling techniques. Recommendations on the use of s-metolachlor in groundwater protection zones will then be formulated as guidelines for policy-makers, administrations and farmers

### Partners

Helmholtz Centre Jülich (DE)

### Financial Support

Administration de la Gestion de l'Eau

### Contact

5, avenue des Hauts-Fourneaux  
L-4362 Esch-sur-Alzette  
phone: +352 275 888 - 1 | [LIST.lu](https://www.list.lu)

© Copyright April 2024 LIST

LUXEMBOURG  
INSTITUTE OF SCIENCE  
AND TECHNOLOGY

